

## A short review on Attention Deficit Hyperactivity Disorder (ADHD): Types, Symptoms, Prevention and Treatment

Devrat Singh<sup>1</sup>, Aman Kumar<sup>2</sup>, Ramandeep Kaur<sup>3</sup>, Suraj Mandal<sup>4\*</sup>, Sanjeev Kumar<sup>5</sup>

<sup>1,2,3,4,5</sup> Pt. Rajendra Prasad Smarak College of Pharmacy, Campus- KajriNiranjanpur, Khutar Road, Puranpur, Pilibhit, Uttar Pradesh, India, 262122

Submitted: 05-02-2022

Accepted: 20-02-2022

### ABSTRACT-

In this present review studying about the attention deficit hyperactivity disorder which may cause sleep disease, anxiety, depression and certain type of learning disabilities etc. Former enquiry command linked meaningful commutability in consideration deficiency/hyperactivity complaint. It is by and large previously analyzed in immaturity and as often as possible keeps going into greater part. Youngsters with ADHD might experience difficulty focusing, controlling rash activities or be excessively action. Attention inadequacy hyperactivity objection (ADHD) is recorded by thought, hyperactivity and impulsivity in compatibility of DSM-IV. ADHD frequency calculation all over the world by and large got across by mythological courses and remedy everywhere the once many ten years hold power enterprises about whether the right diffusion of the complaint has accelerated ceased movement.

**Keywords:** ADHD, DSM-IV, attention deficit hyperactivity disorder.

### I. INTRODUCTION:

ADHD is one of the most common neurodevelopmental diseases of nonage. It is generally first diagnosed in nonage and frequently lasts into majority. Children with ADHD may have trouble paying attention, controlling impulsive actions or be exorbitantly action.<sup>[1]</sup> Attention deficiency hyperactivity complaint (ADHD) is inscribed by consideration, hyperactivity and impulsivity in pursuance of DSM-IV.<sup>[2]</sup> The world Health Organization (WHO) utilizing apart name hyperactive complaint (hyperkinetic disorder) but lists analogous functional yard stick for the complaint. Heed lower of the name used ADHD/HD is one of the most thoroughly delved diseases in drug.<sup>[3, 4]</sup> It has been allied with an expanded range of negatory issues for affected subjects and with a solemn burden to families and society.<sup>[5, 6, 7]</sup> which inscribe it as a

major public health problem understanding of the epidemiological aspect of ADHD/HD may give sapience into its dispensation and origin specifically (etiology) as well as acquaintance for convening the allotment of finances for internal health services in once a period of 10 years especially (decades), in experimenter (inquisitor) from all regions of the world have made stantial sweats.<sup>[8, 9]</sup>

### Types:

there are three different types of ADHD depending on which types of symptoms are strongest in the individual persons:

- 1. Predominantly Inattentive Presentation:** It is difficult for the particular to manage or get over an action. To give consideration to expending or to adhere reference or exchange. The individual is fluently anxious or forgets specific thing of diurnal banality.<sup>[10]</sup>
- 2. Predominantly hyperactive-impulsive presentation:** The individual becomes suddenness and a loquacious person. It is difficult to squat stagnant for prolong. lower children may run jump or climb constantly. The person feels restless and has trouble with impulsivity.
- 3. Combined presentation:** The combined form of above both two have seen in this or inversely present in this.<sup>[11]</sup>

### Symptoms:

Now the first step to know the symptoms start to talk the health care provider to find out the correct symptoms, but every time it has the different symptoms. The different type of ADHD has different symptoms. some of the symptoms are given below:

- Like sleep disease
- Anxiety
- Depression
- Certain type of learning disabilities
- Analogous symptoms
- Day dreams a lot

- Talk too much
- Have difficulty getting along with others<sup>[12]</sup>

On the basis of type of ADHD symptoms are seen if criteria inattention and hyperactivity-impulsivity both were present for last six months then it is combined presentation. If enough symptoms of inattention, but not hyperactivity-impulsivity were present for the past six months then it is predominantly inattentive presentation.<sup>[13]</sup> And the last one is predominantly hyperactive-impulsive presentation it has the enough symptoms of hyperactivity-impulsivity but not intention. The American academy of pediatrics (AAP) recommends that health care providers ask parents, preceptors and other grown-ups who watch for the child about the child's geste in different settings like at home, academy or with peers. The health-care provider should examine that which type of the symptoms does the child have.<sup>[14]</sup>

#### Prevention:

The prevention of ADHD generally precludes comforting or specifics, either single or in admixture. Whileremedy may ameliorate lengthy tour issues, it does not master relief of adversarial issues entirely.<sup>[15]</sup> Specific operated contain instigations, atomoxetine which makes you tired and sleepy, nascence-2 adrenergic receptor agonist, and occasionally antidepressant.<sup>[16,17]</sup> in those who hold condition fastening on lengthy tour prices a big quantum of admiring under pinning worsen duty interpretation.<sup>[18]</sup> ADHDinstigation alike ameliorate continuity and assignment version in kids with ADHD.<sup>[19, 20]</sup>

#### Treatment:

The advice for ADHD treatment means many modes paths involving pharmacotherapy and psychosocial in difficult situation as the major part efficient. earliest range specific for the remedy of ADHD are instigations (methylphenidate, hybrid amphetamine mariners, and amphetamine derivations lisdexamfetamine and dextroamphetamine). Alternate file specifics are atomoxetine, tricyclic antidepressant, bupropion and nascence agonist meta-anatomies and methodical retrospect display moderate to hefty fate sizes of instigation in low hitch discount of symptom.<sup>[21, 22]</sup>

## II. CONCLUSION:

ADHD is a disease which spread most widely in nonage. It is a neurodevelopment disorder characterized by inattention, hyperactivity, and impulsivity, which are extensive impairing and age inappropriate. Due to affect this condition any child cannot be concentrate/attention properly. It is envi-

ronmental as well as a genetic problem. It is usually spread into the child at the age of under twelve years. Some are the very effective symptoms of this disease like children have difficulty in playing and controlling on behavior. It is treated based on the symptoms of the disease. It is treated by the psychotherapy and as well as change in the lifestyle of child. Some medications are given which are stimulant like methylphenidate and mixed amphetamine salts.

**Conflict of Interest:** The authors declare that the review was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest

**Acknowledgement:** The authors are thankful to his/her parents.

**Funding:** None.

**Ethics approval and consent to participate:** Not applicable.

**Human and animal rights:** Not applicable.

**Consent for publication:** Not applicable.

**Code availability:** Not applicable.

## REFERENCE:

- [1]. Jaeschke RR, Sujkowska E, Sowa-Kućma M. Methylphenidate for attention-deficit/hyperactivity disorder in adults: a narrative review. *Psychopharmacology*. 2021 Oct;238(10):2667-91
- [2]. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders DSM 4 TR (Text Revision)*. American Psychiatric Association; 2003.
- [3]. World Health Organization. *The ICD-10 classification of mental and behavioural disorders: clinical descriptions and diagnostic guidelines*. World Health Organization; 1992.
- [4]. Goldman LS, Genel M, Bezman RJ, Slanetz PJ. *Diagnosis and Treatment of Attention-Deficit/Hyperactivity Disorder*. *Jama*. 1998 Apr 8;279(14):14.
- [5]. Dulcan M. Practice parameters for the assessment and treatment of children, adolescents, and adults with attention-deficit/hyperactivity disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*. 1997 Oct 1;36(10):85S-121S.
- [6]. Swanson JM, Sergeant JA, Taylor E, Sonuga-Barke EJ, Jensen PS, Cantwell DP. Attention-deficit hyperactivity disorder and hyperkinetic disorder. *The Lancet*. 1998 Feb 7;351(9100):429-33.

- [7]. Stubbe DE. Attention-deficit/hyperactivity disorder overview: Historical perspective, current controversies, and future directions. *Child and Adolescent Psychiatric Clinics of North America*. 2000 Jul 1;9(3):469-79.
- [8]. Kumawat AR, Singh K, Mangal G. SCOPE OF PANCHAKARMA IN NEURO-DEVELOPMENTAL DISORDERS.
- [9]. Bakare MO. Attention deficit hyperactivity symptoms and disorder (ADHD) among African children: a review of epidemiology and co-morbidities: original. *African journal of psychiatry*. 2012 Sep 1;15(5):358-61.
- [10]. Magnin E, Maurs C. Attention-deficit/hyperactivity disorder during adulthood. *Revue neurologique*. 2017 Jul 1;173(7-8):506-15.
- [11]. Wamulugwa J, Kakooza A, Kitaka SB, Nalugya J, Kaddumukasa M, Moore S, Sajatovic M, Katabira E. Prevalence and associated factors of attention deficit hyperactivity disorder (ADHD) among Ugandan children; a cross-sectional study. *Child and adolescent psychiatry and mental health*. 2017 Dec;11(1):1-7.
- [12]. Daley D, Hollis CP. The World Federation of ADHD international consensus statement: 208 evidence-based conclusions about the disorder.
- [13]. Barbaresi WJ, Colligan RC, Weaver AL, Voigt RG, Killian JM, Katusic SK. Mortality, ADHD, and psychosocial adversity in adults with childhood ADHD: a prospective study. *Pediatrics*. 2013 Apr 1;131(4):637-44.
- [14]. Faraone SV, Banaschewski T, Coghill D, Zheng Y, Biederman J, Bellgrove MA, Newcorn JH, Gignac M, Al Saud NM, Manor I, Rohde LA. The world federation of ADHD international consensus statement: 208 evidence-based conclusions about the disorder. *Neuroscience & Biobehavioral Reviews*. 2021 Sep 1;128:789-818.
- [15]. Shaw M, Hodgkins P, Caci H, Young S, Kahle J, Woods AG, Arnold LE. A systematic review and analysis of long-term outcomes in attention deficit hyperactivity disorder: effects of treatment and non-treatment. *BMC medicine*. 2012 Dec;10(1):1-5.
- [16]. American Psychiatric Association. Diagnostic and statistical manual of mental disorders DMS V. Diagnostic and statistical manual of mental disorders.
- [17]. Bidwell LC, McClernon FJ, Kollins SH. Cognitive enhancers for the treatment of ADHD. *Pharmacology Biochemistry and Behavior*. 2011 Aug 1;99(2):262-74.
- [18]. Modesto-Lowe V, Chaplin M, Soovajian V, Meyer A. Are motivation deficits underestimated in patients with ADHD? A review of the literature. *Postgraduate medicine*. 2013 Jul 1;125(4):47-52.
- [19]. Singletary C. The Effect of Resistance Exercise Circuit Training on Adults with Attention-Deficit/Hyperactivity Disorder (ADHD) (Doctoral dissertation, Southeastern Louisiana University).
- [20]. Modesto-Lowe V, Chaplin M, Soovajian V, Meyer A. Are motivation deficits underestimated in patients with ADHD? A review of the literature. *Postgraduate medicine*. 2013 Jul 1;125(4):47-52.
- [21]. Van der Oord S, Prins PJ, Oosterlaan J, Emmelkamp PM. Efficacy of methylphenidate, psychosocial treatments and their combination in school-aged children with ADHD: a meta-analysis. *Clinical psychology review*. 2008 Jun 1;28(5):783-800.
- [22]. Klassen A, Miller A, Raina P, Lee SK, Olsen L. Attention-deficit hyperactivity disorder in children and youth: a quantitative systematic review of the efficacy of different management strategies. *The Canadian Journal of Psychiatry*. 1999 Dec;44(10):1007-16.