Formulation and Evaluation of Nagarmotha hair oil

Seema Baba Dadas, Gauri Sanjay Maske, Asst.Prof Ranjit Madhukar Shelake, Snehal Satish Deshmane, Pratibha Milind Kore, Sadik Fayaz Shaikh, Rushikesh Shankar Bhosle, Neha Sunil Kamble. Institution Sojar College of Pharmacy, Khandavi, Tal.Barshi, Dist.Solapur.

Date of Submission: 27-05-2023 Date of Acceptance: 08-06-

2023

Abstract: -

Herbal Formulation than Synthetic drug have Less Side effects than synthetic drug so herbal attracted considerable formulation has attention. The largely Chemical Compound of the herbs are essential oils flavonoids, terpenoids, cyperacea. The major chemical components of this herb are essential oils.flavonoids, terpenoids, sesquiterpenes, cyproten, cyperene, aselinene, rotundone, valencene, cyperol, gurjunene, transcalamenenene, cad\'Alene, cyperoutundone, mustskone, isocyperol, acyperone etc. The study investigation of oil isolation from Cyperus rotundus and chemical composition of essential oil of roots of C. Rotundus commonly known as nagarmotha in India in ayurveda C.Rotundus rhizome are considered aromatic, astringent, analgesic, diuretic, carminative stimulant, antitussive, sedative, antibacterial, anti helmintic, antiinflammatory, Anti- rheumatic. The goal of the present study is preparation of & evaluation of poly herbal have hair of using Emblica of cinalis [amia] Alovera. I Botanical Name - Aloe barbadensis miller], Almond Botanical Name -Prunus dulcis rose) Jasmin [Botanical Name-Jasminum [olives]. C.rotandus commonly called as Nagarmotha Purple Nut sedge, or Nut Cyperaceae family. Brass Is a perineal grass that belongs to cyperceae family.

Kevwords: -Hair, Polyherbal, oil, evaluation.

I. Introduction: -

Cyperus. Rotundus-Family: - cypereceae

Humans completely depend on mother nature

as if the primary source of numerous medicines for thousands of years.

In Asian country's C. rotundusrhizomes are used in the treatment of stomach and bowel disorder, hair disorder & inflammatory disease have been widely investigated. There are phytochemical surveys of C.rotundus that contain flavoinds, glycoside, tannis, Monoterpenes, terpenoids, carbohydrate, protein, Separated Amino acid, essential oil, Saponins. Nagarmotha is also called purple nut. sedge or Nut-grass is a common perineal weed with slender, rhizome bulbous at the base & arising singly from the tubers which are about 1-3 cm long. Hair is one of the most important parts of the body derived from ectoderm of theskin. The hair follicle has the unique ability like nail to the regenerate itself.

Structure of hair: -

HAIR STRUCTURE



Hair Growth Cycle: -

The growth and loss of hair like a

DOI: 10.35629/7781-080321922196 | Impact Factor value 7.429 | ISO 9001: 2008 Certified Journal Page 2192



International Journal of Pharmaceutical research and Applications Volume 8, Issue 3, May-June 2023, pp: 2192-2196 www.ijprajournal.com ISSN: 2456-

simple process may seem but the hair growth cycle is actually composed Four distinct phase. This stage of hair growth has been studied in great depth to better understand low hair grows & what might be done to hair loss prevent or treat premature hair loss. The First three

phases &telogen - anagen, catagen cover the growth & matinal. ion of hair & the activity of the hair follicles that produce individual hairs During the final or exogen, phase, "old" hair sheds through usually new hair is getting ready to take its Place [figure]




TABLE 1: VERNACULAR NAMES OF CYPERUS ROTUNDUS

TABLE 1. VERITAGOLAR MARINES OF OTT ERSO ROTORIDOS			
English	Coco grass, Nut grass, Nut sedge, Purple Nutgrass, Purple Nutsedge		
Hindi	Nagarmotha, korehi-jhar, motha, mutha		
Sanskrit	Chakranksha, CharukesaraAbda, Ambuda		
Urdu	Saad kuf		
Arabic	Soad, Soadekufi		
Bengali	Nagarmotha, Moothoo, Musta		
Burma	Vomonniu,		
Gujrat	Nagaramothaya		
Malya	Muahkezamin		

Table 2: Taxonomic Classification of Cyperus Rotundus

Taxonomic Rank	Taxon	
Kingdom	Plantae	
Division	Angiosperms	
Class	Liliopsida	



Volume 8, Issue 3, May-June 2023, pp: 2192-2196 www.ijprajournal.com ISSN: 2456-

Subclass	Commelinids	
Order	Poales	
Family	Cyperaceae	
Genus	Cyperus	
Species	Rotundus	
Common Names	Nutgrass, Nagarmotha	

Benefits of nagarmotha

- 1] Nagarmotha controls hair fall and reduces dandruff.
- 2] Regular use of nagarmotha oil enhances hair texture, adds shine and stimulates hair growth.
- 3] Reduce dandruff and removes excessive dryness.
- 4] Nagarmotha helps to Control hair loss associated with dandruff.
- 5] It also prevents hair fall and premature greying of hair.
- 6] Effective on split ends.
- 7 Repairs damage hair.
- 8] Stimulates hair growth
- 9] Provides Volume to hair.
- 10] Treats and moisturizes dry scalp.

Therapeutic uses of Cyperus Rotundus

Anti-Inflammatory Activity Analgesic Activity Anti-Convulsant Activity Anti-diarrhoeal Activity Anti-Microbial Activity Anti-Cancer Anti-Oxidant

Phytochemistry of Cyperus rotundus: -

rotundus contains several phytochemical constituents of p various classes like fat, gum resins, carbohydrates, essential oils alkaloids, saponins, flavonoids, terpenoids, mono-and sesquiterpenes, albuminous

Steroidal:

glycoside, sitosterol-(6'- hentriacontanoic)-ß-dgalactopyranoside, sitosterol, stigmasterol, sitosterol glucoside, stigmasterol glucoside, chrysoeriol, kaempferol, luteolin, quercetin, rutinand khellol -D-glucopyranoside are known steroids which are present in Cyperus rotundus

Flavonoids: -

Flavonoids like vitexin, orientin, cinaroside, quercetin 3-O-B-D-glucopyranoside, and myrcetin 3-O-B-D-glucopyranoside, isnagin, khellin, ammiol, isorhamnetin, and tricin, luteolin

3-methyl ether, Luteolin 7, 3`-dimethyl ether, Luteolin 5, 3-dimethyl ether, luteolin4-glucoside, Luteolin 7-diglucoside, tricin 5-glucoside, kaempferol are present in Cyperus rotundus. A novel falavonol derivative is discovered in this plant named as cyperaflavoside (myricetin 3,3',5'- trimethyl ether 7-O-B-D-glucopyranoside)

Sesquiterpenes

C. rotundus possess diverse skeletons of sesquiterpenes such patchoulane, rotundane, eudesmane, quaiane, cadinane and caryophyllene types. Patchoulane type sesquiterpenes such as patchoulane-1 (cyperene-3, 8-dione), patchoulane-2 hydroxy cyperoutundone), patchoulane-3 (14acetoxy cyperoutundone), patchoulane-4 (3ßhydroxy- cyperenoic acid), patchoulane-5 (sugetriol-3, 9- diacetate) is present in Cyperus rotundas. Whereas eudesman-6 and 7 are known to be present in this plant. Other eudesmaneare present in this plant. Two novel sesquiterpenes named isocyperotundone (1) and 1,4-epoxy-4-hydroxy-4,5-seco-guain-11-en-5 -one (2) are reported to be present in this plant

II. Methodology: Formulation of Nagarmotha hair oil

The various ingredients used in the formulation of Nagarmotha hair oil. Accurately weight dried roots of Nagarmotha roots and other herbs such as, aloe vera, Amla, almond oil, Coconut oil, Jasmin oil, Cedarwood oil. Where grinded in the mixture and was mixed in 6 ml of almond oil. The above content was boiled for



Volume 8, Issue 3, May-June 2023, pp: 2192-2196 www.ijprajournal.com ISSN: 2456-

15 min. and was filtered through muslin cloth. The filtrate almond oil was added to make up the volume (6 ml). Finally small amount of

colour and flavouring agents was added to the oil and it was placed in Umber coloured bottle.

Ingredients used in Nagarmotha hair oil

Sr. No	Ingredient	Quantity
1	Nagarmotha Powered	10gm
2	Almond oil	12ml
3	Aloe vera Gel	8gm
4	Amla powered	8gm
5	Jasmine oil	4ml
6	Coconut oil	8ml

Formulation and evaluation of nagarmotha hair oil:

1]Sensitivity test:

The formulated nagarmotha hair oil was subjected to the physical and biological evaluation

2]Sensitivity test:

 The prepared Nagarmotha hair oil was applied on 1 cm skin of hand and exposed to sunlight for 4-5 min

3]Acid value:

• Preparation of 0.1 molar solution weighed 0.56g KOH pellets and dissolved In 100 ml of distilled water and stirred continuously. The prepared 0.1 molar KOH solution was filled in a burette preparation of sample: measured 10 ml oil and dissolved in 25 ml of ethanol and 25 ml of ether mixture and shaked. Added 1 ml of phenolphthalein solution and titrated with 0.1 molar KOH solution.

4|Saponification value:

- Accurately weighed 1ml of oil into a 250 ml of conical flask and 10 ml of ethanol: ether mixture was Added to this flask 25ml of 0.5n alcoholic KOH was kept the flask for 30 minutes and the flask was cooled.
- \bullet The cooled solution was titrated against 0.5 N HCL using phenolphthalein indicator similarly the blank titration was performed without

taking oil. Amount of KOH in mg in calculated. 5] **PH:**

The Ph Nagarmotha hair oil was determined using PH meter.

6]Viscosity:

The viscosity was determined using Ostwald's viscometer.

7]Specific gravity:

• Take the specific gravity bottle, rinsed it with distilled water, dry it oven for 15 min, cool, closed it with cap and weight it, now fill the same specific gravity bottle with the sample and closed it with cap and again weigh it. Determine the weight of sample per millimetre by subtracting the weight.

8] Spreadability: -

Oil is spread easily without too much drag and did not produce greater friction in the rubbing process.

9] Oraganoleptic Property:

Oraganoleptic Properties such as colour, odour was determined manually.

Result: -Nagarmotha hair oil improve hair texture, adds shine, stimulates hair growth and provide numerous nutrients. It also prevents hair fall and premature greying of hair. Nagarmotha hair oil prepared from various herbs given in above information.

Sr. No	Param	neter Inference	
1	Sensitivi	ity Test No irritation	
2	Irritatio	on Test No irritation	



Volume 8, Issue 3, May-June 2023, pp: 2192-2196 www.ijprajournal.com ISSN: 2456-

3	Acid value	3.9
4	Grittiness	Smooth
5	PH	7

III. Conclusion:

Cyperus rotundus is one of the important medicinal herbs which is used to treat a number of disease or disorder since ancient times. Nagarmotha is useful for treating dandruff and revitalizing dull, lifeless hair. From the present investigation it was found that the formulated hair oil has optimum standard.

Reference:

- [1]. Gurib-Fakim A: Medicinal plants: traditions of yesterday and drugs of tomorrow. Molecular aspects of Medicine. 2006; 27(1): 1-93.
- [2]. Samuelsson G: Drugs of natural origin: a textbook of pharmacognosy, 5th Swedish Pharmaceutical Press. Stockholm, Sweden 2004.
- [3]. Sri Ranjani Sivapalan
- [4]. Dang GK. Parekar RR. Kamat SK, Scindia AM, Rege NN., Anti-inflammatory activity of Phyllanthus Emblica, Plumbago zeylanica and Cyperus rotundus in acute models of inflammation., Phytother Res.2011Jun;25(6):904-8. doi: 10.1002/ptr.3345. Epub 2010 Dec 3.
- [5]. Gupta MB, Palit TK, Singh N, Bhargava KP. Pharmacological studies to isolate the active constituents from Cyperus rotundus possessing anti-inflammatory, anti-pyretic and analgesic activities. Indian Journal of Medical Research 1971; 59: 76-82.
- [6]. Won-Gil Seo, Hyun-Ock Pae, Gi-Su Oh, Kyu-Yun Chai, Tae-Oh Kwon, Young-Gab Yun, Na-Young Kim, Hun-Taeg Chung. Inhibitory effects of methanol extract of Cyperus rotundus rhizomes on nitric oxide and superoxide productions by murine macrophage cell line, RAW 264.7 cells, Journal of ethnopharmacology, Volume 76, Issue 1, June 2001, Pages 59-64.
- [7]. 6.Miss.Patil Shivani Navnath, Prof.Shinde, R.R. Dr. Hingane L.D. Ijraset Journal for

- Research in Applied Science and Engineering Technology
- [8]. Medically reviewed by Cynthia Cobb, DNP, APRN, WHNP-BC, FAANP - By James Roland on September 25, 2020.
- [9]. Honey J, Neha B. A review on pharmacognosy of Cyperus species. Available from: http://www.pharmatutor.org/articles/
- [10]. Kirtikar KR, Basu BD. Indian Medicinal Plants. 2nd ed., Vol. IV. Dehradun: International Book Distributors; 2007. p. 580-878.
- [11]. Classification of Cyperus rotundus L. United State Department of Agriculture. Available from: http://www.plants.usda.gov/java/
- [12]. Miss.Patil Shivani Navnath ,Prof.Shinde, R.R. Dr.Hingane L.D. Ijraset Journal For Research in Applied Science and Engineering Technology
- [13]. Hikino H, Sunaki N and Takemoto T: Structure and absolute configuration of kusunol. Chemical and Pharmaceutical Bulletin 1968; 16(5): 832-8 83.
- [14]. Kalani PS. SHARMA A, Singh AN, Singh IP and Chhabra BR: Biogenetically important sesquiterpenes from Cyperus notanda. Fitoterapia (Milano). 1995; 66(1).
- [15]. Ohira S, Hasegawa T, Hayashi KL, Hoshino T, Takaoka D and Nozaki H: Sesquiterpenoids from Cyperus rotunda. Phytochemistry 1998, 47(8): 1577-81.
- [16]. Kilani S, Abdelwahed A. Ammar RB, Hayder N. Ghedira K. Chraif L Hammum M and Chekir-Ghedira L Chemical composition, antibacterial and antimutagenic activities of essential oil from (Tunisia) Cyperus tunes. J of Essential Oil Research 2005; 17(6): 695-00
- [17]. Kumar M. Rani M and Meher B Review on Pharmacology and Phytochemistry of Cyperus mundus L Current Research in Pharmaceutical Sciences 2017: 11-5. 87. Kim S, Kim HJ, Kim HI, Jang YP, Oh MS and Jang DS:



Volume 8, Issue 3, May-June 2023, pp: 2192-2196 www.ijprajournal.com ISSN: 2456-

- [18]. Sayed HM, Mohamed MH, Farag SF, Mohamed GA and Proksch P: A new steroid glycoside and furochromones from Cyperus rotundus L. Natural Product Research 2007; 21(4): 343-50.
- [19]. Sayed HM, Mohamed MH, Farag SF and Mohamed GA: Phytochemical and biological investigations of Cyperus rotundus L. Bull Fac Pharm Cairo Univ 2001; 39(3): 195-03.
- [20]. Ibrahim SR, Mohamed GA, Alshali KZ, Haidari RA, El- Kholy AA and Zayed MF: Lipoxygenase inhibitors flavonoids from Cyperus rotundus aerial parts. RevistaBrasileira de Farmacognosia. 2018; 28(3): 320-4.
- [21]. Sayed HM, Mohamed MH, Farag SF and Mohamed GA: Phytochemical and biological studies of Cyperus rotundus L. growing in Egypt. Bull Pharm Sci Cairo Univ 2001; 39: 195-03.
- [22]. Kim SJ, Kim HJ, Kim HJ, Jang YP, Oh MS and Jang DS: New patchoulane-type sesquiterpenes from the rhizomes of Cyperus rotundus. Bulletin of the Korean Chemical Society 2012; 33(9): 3115-8.
- [23]. 88. Xu HB, Ma YB, Huang XY, Geng CA, Wang H, Zhao Y, Yang TH, Chen XL, Yang CY, Zhang XM and Chen JJ: Bioactivity-guided isolation of anti-hepatitis B virus active sesquiterpenoids from the traditional Chinese medicine: Rhizomes of Cyperus rotundus. Journal of Ethnopharmacology 2015; 171: 131-40.
- [24]. Wang Q. Yi C, Duan W, Duan Y, Lou J, Zeng g and yin j: two New Sesquiterpenoids Isolated from Cyperus rotundus L. Natural Product Communications 2021; 16(2): 1934578X21991687.
- [25]. Isha Kumari, Hemlata Kaurav and Gitika Choudhary.
- [26]. Sri Rajan Sivapulan
- [27]. Prof.Dr Ali Esmail Al-Snafi
- [28]. Miss Patil Shivani Navnath, prof Shinde R.R., Dr. Hingane L.D.